

## ICAR-National Research Centre for Integrated Pest Management, Pusa, New Delhi

### Weekly Status Report on Insects Pests & Diseases of Crops

Name of Institute: ICAR - INDIAN INSTITUTE OF SPICES RESEARCH, KOZHIKODE 673 012, KERALA

Date: 11.07.2016 - 17.07.2016

Crop	Crop Stage	Location (with GPS)	Major Insect Pests		Major Plant Diseases		Other Pests (Nematodes, Rat, etc.) (Scientific Name)	Pest Advisories
			Name (Scientific Name)	Status (Low, Medium & Severe)	Name (Scientific Name)	Status (Low, Medium & Severe)		
Black pepper	(a) Vegetative/ Initiation of spikes	Idukki, Kozhikode, Wayanad (Kerala), Kodagu (Karnataka)	<b>Leaf gall thrips</b> ( <i>Liothrips karnyi</i> )	Medium	<b>Stunt disease</b> ( <i>Cucumber mosaic virus, Piper yellow mottle virus</i> )	Low	<b>Nematodes</b> ( <i>Radopholus similis, Meloidogyne incognita</i> ) (Nursery)	<b>Field: Stunt disease</b> Regular monitoring. Remove infected vines and destroy by burning or burying deep in soil. Control the vector (mealy bugs) by drenching with chlorpyrifos (0.075%). <b>Foliar infection due to Phytophthora capsici</b> After the receipt of few monsoon showers, all the vines are to be drenched at a radius of 45-50 cm with copper oxychloride 0.2% @ 5- 10 litres/vine. A foliar spray with Bordeaux mixture 1% is also to be given. Alternatively, drenching and spraying with potassium phosphonate 0.3% @ 5-10 litres/
	(b) Nursery		<b>Top shot borer</b> ( <i>Cydia hemidoxa</i> )	Medium	<b>Foliar infection</b> (due to <i>Phytophthora capsici</i> )	Medium		

					<p><b>Anthracnose</b> (<i>Colletotrichum gloeosporioides</i>) (Nursery)</p> <p><b>Basal wilt</b> (<i>Sclerotium rolfsii</i>) (Nursery)</p> <p><b>Viral infection</b> (Nursery)</p>	<p>Low</p> <p>Low</p> <p>Low to Medium</p>	<p>vine (drench) or potassium phosphonate 0.3% @ 5-10 litres/ vine (drench) also may to be given.</p> <p><b>Anthracnose</b> Prophylactic spraying with Bordeaux mixture (1%) or carbendazim + mancozeb (0.1%).</p> <p><b>Leaf gall thrips</b> Spray dimethoate (0.05%) during emergence of new flushes on young vines.</p> <p><b>Top shot borer</b> Spray quinalphos (0.05%) on tender terminal shoots; repeat spraying at monthly intervals to protect emerging new shoots.</p> <p><b>Nursery:</b> <b>Foliar infection due to <i>Phytophthora capsici</i></b> If foliar infection is noticed, spray Bordeaux mixture (1%) and drench with copper oxychloride (0.2 %). Alternatively, metalaxyl 0.01% (1.25 g/litre) or potassium phosphonate 0.3% (3 ml/litre) could also be used.</p> <p><b>Anthracnose</b> Spray Bordeaux mixture (1%) alternating with carbendazim (0.1%).</p> <p><b>Basal wilt</b> Remove and destroy affected cuttings along with defoliated leaves.</p>
--	--	--	--	--	--	--	--

								<p>After periodic sanitation, the cuttings should be drenched with carbendazim (0.2%) or Bordeaux mixture (1%).</p> <p><b>Viral infections</b> Regular inspection and removal of infected plants. Regular monitoring for insects and spray with dimethoate (0.05%) whenever insect attack is noticed.</p> <p><b>Scale insects</b> Clip off and destroy severely infested leaves. Spray dimethoate (0.1%). Repeat spraying after 21 days to control the infestation completely. (Initiate control measures during early stages of pest infestation).</p> <p><b>Mealy bug</b> Spray dimethoate (0.05%), once infestation is noticed.</p> <p><b>Nematodes</b> Apply carbofuran (0.1%) @ 50 ml/bag.</p>
--	--	--	--	--	--	--	--	---

<b>Cardamom</b>	<b>(a) Vegetative/ Panicle initiation/ Capsule formation</b>	Idukki, Wayanad (Kerala), Kodagu (Karnataka)	<b>Panicle/Shoot borer</b> ( <i>Conogethes punctiferalis</i> )	Low	<b>Leaf blight</b> ( <i>Colletotrichum gloeosporioides</i> )	Low		<b>Field:</b> <b>Panicle/Shoot borer</b> Spray quinalphos (0.075%) coinciding with emergence of panicles and new shoots. <b>Thrips</b> Under Karnataka conditions, spray Fipronil (0.005%) or Spinosad (0.0135%) after undertaking thrashing. Ensure irrigation after thrashing. <b>Leaf blight</b> Maintain optimum shade level by providing 40-60% filtered light. <b>Katte/ Mosaic</b> Prompt inspection of plantation, detection and rouging of virus sources (infected plants/ volunteers) to reduce re-infection. The removed plants may be burnt or buried deep in soil. Removal of natural hosts like <i>Colocasia</i> and <i>Caladium</i> to destroy breeding sites and check population build-up of the vector. <b>Chlorotic streak</b> Prompt inspection of plantation, detection and rouging of virus sources (infected plants/ volunteers) to reduce re-infection. The removed plants may be burnt or buried deep in soil. <b>Azhukal/Capsule rot</b> Thrashing and cleaning of the plant
	<b>(b) Primary seedling nursery</b>		<b>Thrips</b> ( <i>Sciothrips cardamomi</i> )	Low	<b>Katte/Mosaic</b> ( <i>Cardamom mosaic virus</i> )	Medium		

							<p>basin need to be carried out. Regulate thick shade. Prevent water logging by providing adequate drainage. Destroy disease affected portions and plant debris.</p> <p>Prophylactic sprays with Bordeaux mixture (1%). Alternatively, fosetyl-aluminium (0.2%) or potassium phosphonate (0.3%) can be used. Drench plant basin with copper oxychloride (0.2%).</p> <p><b>Primary seedling nursery:</b></p> <p><b>Damping off or seedling rot</b> Prevent water stagnation by providing adequate drainage. Remove and destroy infected/dead seedlings.</p> <p>When initial symptoms are noticed, drench nursery beds with 0.2% copper oxychloride @ 3-5 litres/m<sup>2</sup>. Repeat drenching two to three times at an interval of 15 days.</p>
--	--	--	--	--	--	--	--

<b>Vanilla</b>	<b>Vegetative/ flowering/ bean formation</b>	Karnataka			<b>Premature yellowing and bean shedding</b> ( <i>Colletotrichum vanillae</i> ) <b>Viral diseases</b> ( <i>Bean common mosaic virus</i> , <i>Bean yellow mosaic virus</i> , <i>Cucumber mosaic virus</i> , <i>Cymbidium mosaic virus</i> )	Medium  Medium		<b>Premature yellowing and bean shedding</b> Provide 50% shade in the plantation. Spray carbendazim – mancozeb (0.25%) at 15 – 20 days interval. <b>Viral diseases</b> Regular inspection and removal of infected plants. The removed plants may be burnt or buried deep in soil. Control of vector (aphids) may be undertaken by spraying dimethoate (0.05%).
<b>Ginger</b>	<b>Vegetative</b>	Kerala, Karnataka, Tamil Nadu	<b>Leaf roller</b> ( <i>Udaspes folus</i> )	Low	<b>Soft rot</b> ( <i>P. aphanidermatum</i> and <i>P. myriotylum</i> ) <b>Leaf spot</b> <i>Phyllosticta zingiberi</i>	Low  Low		<b>Soft rot</b> Seed rhizomes are to be selected from disease free gardens. Treat seed rhizomes with mancozeb (0.3%) or metalaxyl mancozeb (0.125%) for 30 minutes before planting. <b>Leaf spot</b> Spray Bordeaux mixture (1%) or mancozeb (0.2%) or carbendazim (0.2%) when the initial symptoms appear. Care should be taken that the spray solution should reach lower surface of the leaves also. <b>Leaf roller</b> Spraying malathion (0.1%) at 21 days intervals.
<b>Turmeric</b>	<b>Vegetative</b>	Kerala, Tamil Nadu, Andhra Pradesh,	<b>Leaf roller</b> ( <i>Udaspes folus</i> )	Low	<b>Rhizome rot</b> ( <i>Pythium aphanidermatum</i> ) <b>Leaf spot</b>	Low  Low		<b>Rhizome rot</b> Treating the seed rhizomes with mancozeb (0.3%) for 30 minutes prior at the time of planting.

		Telangana			<i>(Colletotrichum capsici)</i>			<b>Leaf spot</b> Spray carbendazim or mancozeb (0.2 %) or copper oxychloride (0.2%). <b>Leaf roller</b> Spraying malathion (0.1%) at 21 days intervals.
--	--	-----------	--	--	---------------------------------	--	--	--